



USE OF METEOROLOGICAL MODELS AS INPUT TO REGIONAL AND MESOSCALE AIR QUALITY MODELS — LIMITATIONS AND STRENGTHS

R. A. PIELKE*[†] and M. ULIASZ[‡]

*Department of Atmospheric Science, Colorado State University, Fort Collins, CO 80523, U.S.A. and

[‡]Mission Research Corporation, ASTeR Division, PO Box 466, Fort Collins, CO 80526, U.S.A.

(First received 30 October 1996 and in final form 11 February 1997. Published April 1998)

Abstract—The importance of meteorological variability and uncertainty is described and discussed in the context of dispersion and chemistry of air pollution. Synoptic, mesoscale, and turbulent scales are defined in relation to pollution dilution. Spatial variability effects due, for example, to synoptic baroclinicity, propagating synoptic and mesoscale features, and surface-forced atmospheric circulations are described. Temporal variability resulting from diurnal and seasonal effects are discussed and examples presented. Among the questions addressed is the importance of differential advection relative to horizontal diffusion at different space and time scales. The concept of delayed diffusion is presented. Among the conclusions is that regulating agencies such as the EPA and NPS have generally not taken sufficient advantage of regional and mesoscale meteorological model-generated wind and turbulence fields, nor used the limits on the accuracy of these models to provide an upper limit to the skill of air quality models. Part of this failure is due to the poor communication by scientific researchers, of model capabilities and limits to the agencies and other users of meteorological model output as part of air quality assessments. © 1998 Elsevier Science Ltd. All rights reserved.

Key word index: Dispersion, long-range transport, terrain-induced, mesoscale modeling, boundary layer.

1. INTRODUCTION

The need to accurately describe the wind and turbulence fields for input into air quality assessment models should be obvious (e.g., Physick *et al.*, 1992; Pielke *et al.*, 1983, 1991; Tucker, 1993; Svensson, 1996a; Kallos, 1996; Berkowitz and Fast, 1996; Physick, 1995; Yamada, 1996; Mueller *et al.*, 1996; Millán *et al.*, 1996; Liu and Carroll, 1996; Schayes *et al.*, 1996; Bornstein *et al.*, 1996; Sharan *et al.*, 1996; Nowacki *et al.*, 1996; Banta, 1990). Otherwise, how could the transport, deposition, and dispersion of a chemical effluent and resultant concentration fields and chemical reactions be accurately represented. Sistla *et al.* (1996), for example, show the influence of meteorological uncertainties with respect to ozone control strategies. Al-Wali and Samson (1996) illustrate the sensitivity of the Urban Airshed Model to the placement of boundary layer measurements with respect to the location of emissions. Svensson (1996b) demonstrated that relative errors in meteorological variables simulated in a model may be larger than the uncertainties in chemical rate constants. That paper emphasized the importance of understanding how errors in the meteo-

rological fields affect the results (and conclusions) from the air quality model. Hanna (1992) discusses the effects of data limitations on the ability to improve short-range dispersion models. Kahl (1996) presents a paper on the prediction of trajectory error.

In this paper we discuss this need with particular emphasis on the mesoscale and regional scale. The limitations and strengths of using meteorological models to obtain wind and turbulence information is presented, as well as uncertainties in the atmospheric flow field analyses. These uncertainties will place limits on what accuracy is achieved in estimating chemical concentration fields from air quality models.

2. REVIEW OF MODELING APPROACHES

Meteorological models have been used in a diagnostic form or in prognostic version (Pielke, 1985; Kumar and Russell, 1996; Zannetti, 1990; Ratto *et al.*, 1994). Prognostic models permit nonlinear terms (e.g., advection) to create a flow field that contains information not present in the observational analysis used to initialize the model. Diagnostic models are constrained by available observations, topographic terrain and linear conservation formula (e.g., the conservation of mass) such that they are unable to

* Author to whom correspondence should be addressed.

obtain spatial and temporal structure smaller than the available input data. Hybrid models exist also where a prognostic model ingests observed data and nudges the simulated field towards the measured observations (e.g., see Hoke and Anthes, 1976; Bigler-Engler *et al.*, 1996; Ching and Pleim, 1996). This analysis procedure is referred to as four-dimensional data assimilation (4DDA) (Fast, 1995). Hariharan and Venkatram (1996) illustrate errors that result when mass conservation is not retained in air quality models. McNider *et al.* (1996) demonstrate the values of 4DDA even when a reasonably complete prognostic model formulation is used.

This paper focuses on modeling of atmospheric flow and turbulence using prognostic models. It should be pointed out that turbulence characteristics required by air pollution models (i.e., eddy diffusivities, wind velocity variances, Lagrangian time scales) are usually not provided directly by meteorological models but must be derived from their output using wind and temperature fields and often some additional fields such as turbulent kinetic energy, turbulent length scales, mixing height or parameters of the atmospheric surface layer. Ideally, the same turbulence parameterization scheme should be used for this purpose as one applied in the meteorological model, otherwise, these calculations may introduce additional sources of errors. The turbulence parameterizations used in recent mesoscale and regional meteorological models are reviewed by Uliasz (1994).

Contemporary air quality models taking into account atmospheric chemistry and removal processes may also require additional meteorological input including radiative and/or water fields (i.e. water vapor, cloud, and precipitation fields). The current state and future directions of tropospheric chemistry and transport models are discussed by Peters *et al.* (1995).

In most modeling approaches atmospheric chemistry and dynamics are effectively decoupled allowing one to use an off-line approach where the meteorological model is applied independently from the air quality model. It is possible to simulate different emission scenarios or to run different dispersion models using the same meteorological fields without necessity to repeat the meteorological simulations. However, the output from meteorological models must be stored for this purpose quite frequently. A modeling domain of the mesoscale/regional meteorological models is often much larger than the domain used in air pollution simulations and it may be covered by multiple nested grids. As a consequence, a huge volume of data must be produced in the case of long-term simulations. Therefore, an additional processing of meteorological fields may be useful to extract only these fields from the meteorological output which are required by the air pollution model and to reduce size of the modeling domain. An example of this procedure is described by Uliasz *et al.* (1996) for the 1 year simulation of air pollution transport in the southwestern United States.

The on-line modeling approach must be used in order to investigate feedback effects between air chemistry and atmospheric dynamics. Therefore, more advanced modeling systems must contain both chemical and meteorological components and allow interaction between the two.

In addition to continuous meteorological simulations, an alternative approach may be considered for practical applications as proposed by Enger (1990). First, a data base with a thousand or more wind and turbulence fields is created by running a prognostic mesoscale model for a typical diurnal cycle with different directions and speeds of the geostrophic wind. To simulate the dispersion, the data base is searched for the meteorological fields which are the closest to the actual observations in the area and these fields are used as input to the dispersion model. This approach was applied in a moderately complex terrain for both operational and long-term dispersion simulations.

3. METEOROLOGICAL VARIABILITY

The dispersion of chemical effluent from point, line and area sources is the result of differential wind advection and turbulent diffusion. Differential wind flow results from time- and space-varying winds, which in the context of a numerical model or observational analyses are those winds that are resolved. The winds are three-dimensional including horizontal and vertical components. The term turbulence is used to describe those scales of flow that are smaller than the model or analysis resolution. As discussed by Pielke (1991), models are able to resolve features reasonably only when the spatial scale is at least four grid increments in each of the three directions.

The resolved scales can be separated into two categories — the synoptic scale where the flow is in near gradient wind balance above the planetary boundary layer and the mesoscale where the flow deviates significantly from the balance even in the free atmosphere. Both synoptic- and mesoscales are close to hydrostatic balance. Turbulent flows can be considered as deviating significantly from hydrostatic balance; thus, cumulus clouds including thunderstorms are one type of turbulence.

Dilution of chemical effluent occurs due to turbulent diffusion, and because of wet and dry deposition to the ground. This dilution reduces the chemical concentration of a chemical species in the absence of additional effluent or chemical reaction (e.g., due to photochemical effects) which creates more of the chemical than is being diluted. Differential advection, in the absence of deposition and chemical reactions, can only result in the vertical and horizontal displacement of a chemical but will not result in dilution until turbulent diffusion occurs. If there is a time lag between differential advection and when diffusion

occurs, this is referred to as delayed dispersion (Moran, 1992).

3.1. Spatial effects

3.1.1. *Synoptic and larger scales.* Meteorological variability on this scale is directly associated with baroclinic effects (i.e., vertical shear of the horizontal wind), propagating synoptic features, and large-scale ascent/descent patterns. Bluestein (1992) and Carlson (1991) provide recent texts on the dynamics of these synoptic effects.

On the synoptic scale, vertical wind shear is a direct result of a horizontal temperature gradient. Since such gradients are a feature of the polar front and extratropical cyclones (e.g., see Pielke *et al.*, 1987), this mechanism of differential wind transport will always

occur near these weather features. The movement of these weather features is what creates ascending and descending air with magnitudes up to tens of centimeters per second over thousands of square kilometer areas. Figure 1 shows how even a horizontal synoptic wind field differential wind flow can deform an area of pollution.

Examples of papers that discuss this effort are Artz *et al.* (1985), Stocker *et al.* (1990).

3.1.2. *Propagating mesoscale systems.* Squall lines, frontal circulations, cumulus convection embedded in stratiform systems and mesoscale convective systems are examples of atmospheric features of hundreds of kilometers in scale that move through the atmosphere. When deep cumulonimbus occurs with these features, rapid and extensive, horizontal and vertical




Fig. 1. Horizontal deformation of a large-scale parcel by two-dimensional incompressible flow at 500mb as predicted by a barotropic numerical model after a time period of: (a) 0 h; (b) 6 h; (c) 12 h; (d) 24 h; and (e) 36 h. The initial streamline pattern is shown at the top. The sides of the colored grid squares are 300 km (Welander, 1955; as reproduced by Moran, 1992).




Fig. 2. Sequential vertical profiles (height above the ground) of the east–west (upper panel) and north–south (lower panel) velocity component winds from 14 balloon launches over a 6 h period on 31 March 1976 at St. Cloud, Minnesota (from Gage and Jaspersen, 1979; as reproduced by Moran, 1992). The profiles on the right-hand side of the figure are collapsed on top of each other on the left side, to better illustrate the temporal variability. Positive values correspond to westerly and southerly components, respectively.

differential winds and turbulent diffusion will result in considerable dilution. Scavenging of aerosols and water soluble chemical species and resultant wet deposition can further dilute the chemical concentrations. In addition to scavenging, deep cumulus convective clouds vertically transport chemical species from near the surface to higher altitudes (Cotton *et al.*, 1995; Lyons *et al.*, 1986).

3.1.3. *Mesoscale surface effects.* Sea and land breezes, mountain–valley winds, and winds generated due to other types of land surface variability (e.g., urban–rural differences, snow–no snow cover variability, vegetated areas adjacent to less vegetated regions, etc.) also contribute to differential wind and turbulence fields. These mesoscale features develop due to horizontal surface gradients in surface sensible and latent turbulent heat fluxes. These atmospheric features are described in a wide variety of sources including (Atkinson, 1981, and Pielke (1984). Avissar (1996) illustrated the importance of vegetation in the urban environment on the dispersion of pollutants in which turbulent mixing is strongly influenced by the relative partitioning of turbulence into sensible and latent fluxes by the plants, as contrasted with non-vegetated surfaces.

Even if the differences in surface turbulent fluxes are not strong enough to create a distinct mesoscale circulation, they can significantly affect vertical structure of the atmospheric boundary layer and in turn atmospheric dispersion. These effects were demonstrated by simulating atmospheric dispersion over a series of land patches with different soil water content (Pielke and Uliasz, 1993).

3.2. Temporal effects

The variability of wind speed and direction, turbulence intensity and precipitation rates will also influence the dilution of chemicals. Even in the free atmosphere, winds are not steady but undergo variations on all time scales (Fig. 2).* McNider *et al.* (1988) shows how inertial oscillations due to the Coriolis effect, synoptic vertical shear, and decoupling and recoupling of the planetary boundary layer with the volume of the chemical species of interest will

* Figure 2 indicates that the routine release of radiosonde balloons at 12 h intervals by the operational weather centers around the world, introduces an appreciable error in the ability to account for dispersion due to time changes in wind speed and direction.




Fig. 3. Instantaneous depiction of plume release after 42 h from Mt. Isa smelter in Australia. (a) Only planetary boundary layer (PBL) turbulence but no PBL shear and no Coriolis force. (b) Including Coriolis force but no PBL turbulence. (c) Includes PBL turbulence, PBL shear and Coriolis force. (d) Same as (c) except for baroclinic case which includes geostrophic shear. The data correspond to particles released from the smelter stack 300 m altitude at 600 s intervals starting at 1115 Local Standard Time. The low-level wind was from right to left (easterly) in the southern hemisphere. The “solid” appearing line is actually the superposition of particles which remain close to each other (from McNider *et al.*, 1988; see that reference for more details on this experiment).




Fig. 4. Hodographs of the summertime diurnal wind variation for three levels at Fort Worth, Texas including the mean wind vector and the wind vectors at the two standard synoptic observing times (0000–1200 UTC). Speeds indicated by the concentric circles are in meters per second. The size of the solid circles for each observing time (shown as CST) is proportional to the magnitude of the probable error in determining the deviation vector at that time (adapted from Bonner and Paegle, 1970 and reported in Moran, 1992).

directly influence dispersion (Fig. 3). Figure 4 shows how the inertial wind variation in the U.S. Great Plains creates a time-varying horizontal wind, as well

as what is missed because the routine synoptic observations are only made at 12 h intervals.

On the longer time scale, seasonal differences in the structure of synoptic and mesoscale systems, in surface-based turbulence, and precipitation type and intensity will also result in different magnitudes of dilution.

4. EXAMPLES OF UNCERTAINTY

4.1. Due to spatial resolution

Increasing spatial resolution of a meteorological model allows one to include more mesoscale motions in its numerical solution. The higher resolution meteorological fields used in the dispersion model with corresponding resolution will in turn lead to lower concentrations simulated on a regional scale, i.e., a given receptor will experience lower contribution from distant emission sources. This effect is illustrated in Fig. 5 on an example of passive tracer transport between the Los Angeles basin and the Spirit Mountain receptor (located about 350 km NE from Los Angeles) (Uliasz *et al.*, 1996). The meteorological simulations were performed with the Colorado State University Regional Atmospheric Modeling System (CSU RAMS) using two nested grids with spacing 60 and 12 km, respectively. Dispersion was simulated by Lagrangian Particle Dispersion model using the output from both RAMS grids (RAMS) and the output from the coarse grid only (RAMS-1). The output from the coarse grid # 1 does not really represent an independent coarse resolution simulation since there is a two-way interaction between nested grids in RAMS. Nevertheless, concentration plots in Fig. 5 demonstrate an evident and important difference between these two sets of particle simulations. The coarse grid particle simulations provide much higher concentration values in comparison to the simulations where input from both nested grids was used. Mesoscale motions simulated on the finer grid # 2 cause much more intensive dispersion of the tracer. This figure shows also that concentrations simulated by the ATAD (Atmospheric Transport and Diffusion) model are higher than those predicted by the RAMS-LPD simulations which take into account more mesoscale motions. The ATAD model is a tool commonly used by the National Park Service (NPS) for a trajectory analysis. In this model, a 2-D wind field (u, v) is interpolated from rawinsonde and pibal data. The wind field is averaged within a transport layer determined from a temperature inversion height.

There are a number of papers that illustrate how the spatial resolution in a model influences how realistically transport is represented. For example, Fig. 6 from Lyons *et al.* (1995) shows how vertical motion associated with a sea breeze at Kennedy Space Center is not accurately represented until the horizontal grid intervals are reduced to 1 km. Banta *et al.* (1996)




Fig. 5. Concentration of a passive tracer at Spirit Mountain during a three-month period (July–August) as simulated for the Los Angeles source by two versions of RAMS/LPD simulations and an ATAD model analysis (concentrations presented at the same scale).




Fig. 6. Maximum model-predicted upward vertical motions w (cm s^{-1}) in the sea breeze convergence zones around the Kennedy Space Center region, 7 November 1988, as a function of the size of the model grid cell (9 km; a two nested run with the smallest grid increment of 3 km; a three model run with the smallest grid increment of 1 km; from Lyons *et al.*, 1995).

discusses the implications of small-scale flow features to modeling dispersion over complex terrain. Portela and Castro (1996) illustrate important (in the context of dispersion) atmospheric features that are missed in synoptic analyses and forecasts from the European Centre for Medium-Range Weather Forecasts (ECMWF) over the Iberian peninsula. Byun (1996) illustrates the influence of the number of vertical levels in the Mesoscale Model Generation 4 (MM4) meteorological model on planetary boundary layer height.

4.2. Due to model type

Mesoscale and regional models have different numerical formulations and parameterization schemes. Pielke and Pearce (1994) recently report on several mesoscale models and their capability in two case study simulations. Schlünzen (1994) overviews several German mesoscale models. Eastman *et al.* (1995)

compares a prognostic model with an EPA-approved diagnostic model for the western shore of Lake Michigan (Fig. 7). The observed tracer data agreed much more with the simulated tracer behavior using the prognostic model. Figure 8 illustrates how the additional physics in the prognostic model resulted in a more realistic behavior of the simulated tracer movement. Lagoiwardos *et al.* (1996) contrasts two prognostic models over Greece. Moran and Pielke (1996a, b) contrast a prognostic model with several diagnostic models for a tracer release from Norman, Oklahoma (see Fig. 9). Grossi *et al.* (1996) recommends the use of output from different meteorological models to assess the sensitivity and the robustness of simulated air quality concentrations.

Several research groups have utilized wind fields from the NGM model to run their dispersion or chemistry models, e.g. Schichtel and Husar (1994); Venkatram *et al.* (1994). The NGM model is a

LMOS TRACER STUDY #2-July 16, 1991 14:00-21:00 CDT
 DRIVE TRACK/SF6 PLUME LOCATION-<500PPT- WEST VIEW






Fig. 7. Perspective view of (a) observations from a mobile van from 1300 to 2000 LST. Values are in ppt (Wilkerson, 1991). (b) Surface isopleths in log (ppt). Left (ISCSST); right (LPDM). The two panels taken at 1600 LST. Domain size is 37.5 km x 37.5 km (from Eastman *et al.*, 1995).

hydrostatic primitive equation hemispheric model used operationally by the National Weather Service. Availability of archived NGM output is very attractive since it allows one to perform a multi-year transport study when combined with a rather simple dispersion model. However, these meteorological fields have low time and space resolution: $\Delta x \approx 80$ km (archived with $2\Delta x$ spacing), vertical grid spacing: $\Delta z \approx 150$ m close to the surface. Therefore, the representation of terrain topography is very crude.

Regional transport simulations in the southwestern United States have been validated using a tracer of opportunity—methylchloroform released mostly in the Los Angeles area and measured in three receptors in the vicinity of the Grand Canyon National Park (Uliasz *et al.*, 1996). While correlations between observed and simulated methylchloroform concentrations are within the range of 0.3–0.8 for different months of 1992 for the results from RAMS/LPD modeling, the correlations for results from dispersion




Fig. 8. Typical plume trajectories, calculated from the prognostic model output by utilizing only surface layer winds (trajectory A) which stay at 30 m altitude and the complete 3D wind field (trajectory B) which rises to heights of 1600 m (from Lyons *et al.*, 1995).




Fig. 9. Schematic showing perfluoromonethylcyclohexane (PMCH) cloud centers (indicated by solid black dots) and approximate cloud widths (indicated by bars passing through cloud centers) predicted by eight mesoscale atmospheric dispersion models for the first 3 h sampling period of the Great Plains experiment (0800–1100 UTC 9 July 1980). Note that RADM corresponds to the radom-walk advection and dispersion model. The Colorado State University modeling system result corresponds to the baseline experiments (experiment 4b). An estimate of the actual tracer puff position is also shown based on an extrapolation of 600 km arc sampler measurements since it is impossible to tell from the measurements when the tracer material first arrived at this arc. The 3 h release of perfluorocarbon tracer took place at Norman, Oklahoma (lower left-hand corner) beginning at 1900 UTC 8 July 1980 (from Moran and Pielke, 1996b).

models driven by NGM output are always negative (Uliasz *et al.*, 1994; Schichtel and Husar, 1994). It seems that the NGM model is not able to reproduce

summer monsoonal flow in southern California due to a lack of sufficient initialization data in this area and coarse resolution. It is interesting that the same

NMC analysis data were used to initialize the RAMS simulations which were much more successful due to better resolution and physics.

4.3. Due to averaging time of the meteorological model simulated fields

It is often assumed that longer averaging time of the model output of winds and turbulence will necessarily reduce the error and uncertainty of the dispersion field. However, this is only true if the shorter-term variability were random. If there are systematic features of the atmospheric flow (e.g., surface-forced mesoscale flow, etc.) that are not resolved by the meteorological model, averaging of the simulated data will be unable to properly represent atmospheric dispersion.

Figure 10 compares observed and simulated methylchloroform concentrations in the southwestern United States with the aid of spectral analysis (Uliasz *et al.*, 1996). The presented coherence is derived from a cospectrum calculated for two time series and varies between 0 and 1 (e.g. Sirois *et al.*, 1995). It may be interpreted as a correlation between these time series expressed as a function of frequency or wavelength. The simulated concentrations at three receptors Spirit Mountain, Meadview and Long Mesa in the vicinity of the Grand Canyon National Park were obtained from RAMS/LPD modeling as a contribution from the Los Angeles basin emission. The coherence between the simulated and observed time series of methylchloroform concentration varies significantly with wavelength (time scale) but have similar features for each receptor site. All cospectra show that simulated time series are to some extent coherent with the observed ones for the time scale longer than about

3 days. For the shorter time scale, the cospectra become very noisy. The maximum coherence (between 0.6 and 0.8) appears at time scales between 150 and 180 h at Spirit Mountain and Meadview. This corresponds to the weekly cycles in methylchloroform concentrations related to the fact that this anthropogenic emission is shut down during weekends. It should be noted that the maximum value of coherence for Meadview is lower than that of the two other sites. The Meadview receptor is located at a mouth of Grand Canyon in a much more complex terrain than other receptors. The horizontal grid spacing of the CSU RAMS in these long-term simulations performed over the entire year of 1992 was 12 km on a finer grid. This spatial resolution is obviously not sufficient to correctly represent terrain features in both source and receptor areas. Although it was possible to correctly simulate long-term episodes (3 days) of the observed methylchloroform, the much higher resolution of meteorological fields would be necessary to reproduce shorter-term episodes strongly affected by local and mesoscale circulations in complex terrain.

5. WHAT NEEDS TO BE DONE

There are opportunities to improve the accuracy of differential wind and diffusion for use in air quality models. These include the use of research and operational meteorological models. For example, since the United States National Weather Service runs an Eta model twice daily every day, why not use the output of that model (archived at half-hourly time periods) to input to a dispersion model?

There are also a number of questions that need to be answered. How can model output (which is a grid-volume representation) be most accurately compared with observed data (which is usually point data)? What effect does ensemble average parameterization representations for turbulence and other physical processes in meteorological models influence the limitation on estimating natural uncertainty? What value-added would be achieved, and what uncertainty range would result from ensemble regional and mesoscale simulations for use as input into air quality models?

Can dispersion models fully utilize and take advantage of information provided by a mesoscale or regional meteorological model? Lagrangian particle models can easily handle meteorological fields of any resolution as input. Lagrangian puff models may have problems with more complicated wind fields (puff splitting techniques try to overcome some difficulties). Eulerian grid models usually have quite poor resolution, especially, in the vertical.

Also, what is the value of advanced meteorological modeling when there is a huge uncertainty in emission data? It is a problem not only with uncertain emission rates but also with uncertain parameters characterizing stack emission. The effect related to uncertainty in




Fig. 10. Cospectra of the observed methylchloroform concentrations and simulated Los Angeles contribution for three receptors (from Uliasz *et al.*, 1996).

plume rise (effective stack height) calculations may be enhanced when meteorological fields with a detailed vertical structure of the atmospheric boundary layer are used in regional dispersion simulations instead of a mixing layer concept.

The level of accuracy imposed on meteorological models by computer and data limitations needs to be explored. For example, model resolution is constrained by computer memory, while initialization and boundary condition information limits model skill. There is a need, for example, for better land cover and soil moisture data.

6. SUMMARY

In this paper we have sought to address the value of meteorological models as input to regional and meso-scale air quality models. Unfortunately, up to the present, regulating agencies such as the U.S. EPA and National Park Service have generally failed to take full advantage of meteorological model-generated wind and turbulence fields, nor used the limits on the accuracy of these models to provide an upper limit to the skill of air quality models. Work is underway in this direction by the EPA (Dennis *et al.*, 1996) and this effort should be strongly supported by the air quality community. Moreover, the EPA minimum recommended performance evaluation procedures (EPA, 1991) need to be extended to meteorological model evaluation as suggested, for instance, by Wheeler (1996).

Acknowledgments—This paper is an outgrowth of an invited presentation at the 5th International Conference on Atmospheric Sciences and Applications to Air Quality held in Seattle, Washington, 18–20 June 1996. The paper was ably prepared and edited by Dallas McDonald. Partial support for this paper was provided by Grant #UAH SUB93-183 from the EPA through the University of Alabama. The authors would also like to thank the Referees for their very useful comments and suggested changes and additions to the paper. Discussions over the years with Mike Moran, Dick McNider, Walt Lyons, Bill Physick, John Garratt, Joe Eastman, and Xubin Zeng were very valuable in developing our perspective on the subject of our paper. Also, Chapter 2 in Mike Moran's Ph.D. Dissertation provides an excellent, comprehensive discussion of atmospheric dispersion.

REFERENCES

Atkinson, B. W. (1981) *Mesoscale Atmospheric Circulations*. Academic Press, New York, 495 pp.

Al-Wali, K. I. and Samson, P. J. (1996) Preliminary sensitivity analysis of urban airshed model simulations to temporal and spatial availability of boundary layer wind measurements. *Atmospheric Environment* **30**, 2027–2042.

Artz, R., Pielke, R. A. and Galloway, J. (1985) Comparison of the ARL/ATAD constant level and the NCAR isentropic trajectory analyses for selected case studies. *Atmospheric Environment* **19**, 47–63.

Avissar, R. (1996) Potential effects of vegetation on the urban thermal environment. *Atmospheric Environment* **30**, 437–448.

Banta, R. M. (1990) The role of mountain flows in making clouds. In *Atmospheric Processes over Complex Terrain*, ed. Blumen, W., Meteorological Monographs, Vol **23**, 229–283.

Banta, R. M., Olivier, L. D., Gudiksen, P. H. and Lange, R. (1996) Implications of small-scale flow features to modeling dispersion over complex terrain. *Journal of Applied Meteorology* **35**, 330–342.

Berkowitz, C. M. and Fast, J. D. (1996) The influence of regional-scale atmospheric circulations on chemical mixing over the western north Atlantic. *Preprints, 9th Joint Conference on Applications of Air Pollution Meteorology with A&WMA*, 28 January–2 February 1996, Atlanta, Georgia, American Meteorological Society, Boston, pp. 117–121.

Bigler-Engler, V., Brown, H. and Wagner, K. K. (1996) A hybrid modeling technique to address coastal meteorology and complex terrain in the San Diego photochemical modeling domain. *Preprints, 9th Joint Conference on Applications of Air Pollution Meteorology with A&WMA*, 28 January–2 February 1996, Atlanta, Georgia, American Meteorological Society, Boston, pp. 429–433.

Bluestein, H. (1992) *Synoptic-Dynamic Meteorology in Midlatitudes*. 431 pp. Oxford University Press, New York.

Bonner, W. D. and Paegle, J. (1970) Diurnal variations in boundary layer winds over the south central United States in summer. *Monthly Weather Review* **98**, 735–744.

Bornstein, R., Thunis, P., Grossi, P. and Schayes, G. (1996) Topographic vorticity-model mesoscale- β (TVM) model. Part II: Evaluation. *Journal of Applied Meteorology* **35**, 1824–1834.

Byun, D. W. (1996) Effects of vertical resolution of a regional photochemical model on the diurnal ozone concentrations in the planetary boundary layer. *Preprints, 9th Joint Conference on Applications of Air Pollution Meteorology with A&WMA*, 28 January–2 February 1996, Atlanta, Georgia, American Meteorological Society, Boston, pp. 203–206.

Carlson, T. N. (1991) *Mid-Latitude Weather Systems*, 507 pp. Harper Collins, London, UK.

Ching, J. K. S., and Pleim, J. E. (1996) Study of gridded mixing heights and cloud fields derived from the mesoscale meteorological model with four dimensional data assimilation. *Preprints, 9th Joint Conference on Applications of Air Pollution Meteorology with A&WMA*, 28 January–2 February 1996, Atlanta, Georgia, American Meteorological Society, Boston, pp. 508–511.

Cotton, W. R., Alexander, G. D., Hertenstein, R., Walko, R. L., McAnelly, R. L. and Nicholls, M. E. (1995) Cloud venting. *Earth Science Review* **39**, 169–206.

Dennis, R. L., Byun, D. W., Novak, J. H., Galluppi, K. J., Coats, C. J. and Vouk, M. A. (1996) The next generation of integrated air quality modeling: EPA's models-3. *Atmospheric Environment* **30**, 1925–1938.

Eastman, J. L., Pielke, R. A. and Lyons, W. A. (1995) Comparison of lake-breeze model simulations with tracer data. *Journal of Applied Meteorology* **34**, 1398–1418.

Enger, L. (1990) Simulation of dispersion in moderately complex terrain — Part C. A dispersion model for operational use. *Atmospheric Environment* **24A**, 2457–2471.

EPA (1991) Guideline for the Regulatory Application of the Urban Airshed Model. Report EPA-450/4-91-013, U.S. Environmental Protection Agency, Research Triangle Park, NC.

Fast, J. D. (1995) Mesoscale modeling and four-dimensional data assimilation in areas of highly complex terrain. *Journal of Applied Meteorology* **34**, 2762–2782.

Gage, K. S. and Jaspersion, W. H. (1979) Mesoscale wind variability below 5 km as revealed by sequential high-resolution wind soundings. *Monthly Weather Review*, **107**, 77–86.

- Grossi, P., Giovannoni, J.-M. and Russell, A. G. (1996) Intercomparison of meteorological models applied to the Athens area and the effect on photochemical pollutant predictions, *Journal of Applied Meteorology* **35**, 993–1008.
- Hanna, S. R. (1992) Effects of data limitations on hopes for improved short range atmospheric dispersion models. In *Proceedings, Objectives for Next Generation of Practical Short-Range Atmospheric Dispersion Models*, (edited by Olesen and Mikkelsen), 6–8 May 1992, Risø, Denmark, pp. 77–85.
- Hariharan, R. and Venkatram, A. (1996) The sensitivity of numerical advection schemes to mass inconsistency in wind fields. *Preprints, 9th Joint Conference on Applications of Air Pollution Meteorology with A&WMA*, 28 January–2 February 1996, Atlanta, Georgia, American Meteorological Society, Boston, pp. 102–106.
- Hoke, J. E. and Anthes, R. A. (1976) The initialization of numerical models by a dynamic-initialization technique. *Monthly Weather Review* **104**, 1551–1556.
- Kahl, J. D. W. (1996) On the prediction of trajectory model error. *Atmospheric Environment* **17**, 2945–2957.
- Kallos, G. (1996) Transport and transformation of air pollutants from Europe to the East Mediterranean Region (T-TRAPEM). Environmental Research programme: Avicenne. Final Report, Contract No. AVI*-CT92-0005, April 1996, Commission of the European Communities, 352 pp.
- Kumar, N. and Russell, A. G. (1996) Comparing prognostic and diagnostic meteorological fields and their impacts on photochemical air quality modeling. *Atmospheric Environment* **30**, 1989–2010.
- Lagoivarados, K., Kotroni, V., Dobricic, S., Nickovic, S. and Kallos, G. (1996) On the storm of 21–22 October 1994 over Greece: Observations and model results. *Journal of Geophysical Research*, in press.
- Liu, M. and Carroll, J. J. (1996) A high-resolution air pollution model suitable for dispersion studies in complex terrain. *Monthly Weather Review* **124**, 2396–2409.
- Lyons, W. S., Calby, R. H. and Keen, C. S. (1986) The impact of mesoscale convective systems on regional visibility and oxidant distribution during persistent elevated pollution episodes. *Journal of Climate and Applied Meteorology* **25**, 1518–1531.
- Lyons, W. A., Pielke, R. A., Tremback, C. J., Walko, R. L., Moon, D. A. and Keen, C. S. (1995) Modeling the impacts of mesoscale vertical motions upon coastal zone air pollution dispersion. *Atmospheric Environment* **29**, 283–301.
- McNider, R. T., Moran, M. D. and Pielke, R. A. (1988) Influence of diurnal and inertial boundary layer oscillations on long-range dispersion. *Atmospheric Environment* **22**, 2445–2462.
- McNider, R. T., Singh, M. P. and Gupta, S. (1996) Nocturnal wind structure and plume growth rates due to inertial oscillation. *Preprints, 9th Joint Conference on Applications of Air Pollution Meteorology with A&WMA*, 28 January–2 February 1996, Atlanta, Georgia, American Meteorological Society, Boston, pp. 455–458.
- Millán, M., Salador, R. and Mantilla, E. (1996) Mesoscale processes and photo-oxidant cycles on the Spanish Mediterranean coast. *Preprints, 9th Joint Conference on Applications of Air Pollution Meteorology with A&WMA*, 28 January–2 February 1996, Atlanta, Georgia, American Meteorological Society, Boston, pp. 434–437.
- Moran, M. D. (1992) Numerical modelling of mesoscale atmospheric dispersion. Ph. D. dissertation, Department of Atmospheric Science, Colorado State University, p. 758.
- Moran, M. D. and Pielke, R. A. (1996a) Evaluation of a mesoscale atmospheric dispersion modeling system with observations from the 1980 great plains mesoscale tracer field experiment. Part I: data sets and meteorological simulations. *Journal of Applied Meteorology* **35**, 281–307.
- Moran, M. D. and Pielke, R. A. (1996b) Evaluation of a mesoscale atmospheric dispersion modeling system with observations from the 1980 Great Plains mesoscale tracer field experiment. Part II: dispersion simulations. *Journal of Applied Meteorology* **35**, 308–329.
- Mueller, S. F., Song, A., Norris, W. B., Gupta, S. and McNider, R. T. (1996) Modeling pollutant transport during high-ozone episodes in the southern Appalachian Mountains. *Journal of Applied Meteorology* **35**, 2105–2120.
- Nowacki, P., Samson, P. J. and Sillman, S. (1996) Sensitivity of Urban Airshed model (UAM-IV) calculated air pollutant concentrations to the vertical diffusion parameterization during convective meteorological situations. *Journal of Applied Meteorology* **35**, 1790–1803.
- Peters, L. K., Berkowitz, C. M., Carmichael, G. R., Easter, R. C., Fairweather, G., Ghan, S. J., Hales, J. M., Leung, L. R., Pennel, W. R., Potra, F. A., Saylor, R. D. and Tsang, T. T. (1995) The current state and future direction of Eulerian models in simulating the tropospheric chemistry and transport of trace species: a review. *Atmospheric Environment* **29**, 189–222.
- Physick, W. L. (1995) Photochemical smog studies in Australian cities. In *Urban Air pollution, Vol. 2* (edited by Poer, H. and Moussiopoulos, N). Computational Mechanics Publications, Southampton, Boston.
- Physick, W. L., Noonan, J. A., Manins, P. C., Hurley, P. J. and Malfroy, H. (1992) Application of coupled prognostic windfield and Lagrangian dispersion models for air quality purposes in a region of coastal terrain. In *Air Pollution Modeling and Its Application IX*, eds. H. Van Dop and G. Kallos, pp. 725–729, Plenum Press, New York.
- Pielke, R. A. (1984) *Mesoscale Meteorological Modeling*. 612 pp. Academic Press, New York.
- Pielke, R. A. (1985) The use of mesoscale numerical models to assess wind distribution and boundary layer structure in complex terrain. *Boundary-Layer Meteorology* **31**, 217–231.
- Pielke, R. A. (1991) A recommended specific definition of “resolution”. *Bull. Amer. Meteor. Soc.* **72**, 1914.
- Pielke, R. A. and Pearce, R. P. (1994) *Mesoscale Modeling of the Atmosphere*. American Meteorological Society Monograph, Vol. 25, 167 pp.
- Pielke, R. A., McNider, R. T., Segal, M. and Mahrer, Y. (1983) The use of a mesoscale numerical model for evaluations of pollutant transport and diffusion in coastal regions and over irregular terrain. *Bulletin of the American Meteorological Society* **64**, 243–249.
- Pielke, R. A., Garstang, M., Lindsey, C. and Gusdorf, J. (1987) Use of a synoptic classification scheme to define seasons. *Theoretical Applied Climatology* **38**, 57–68.
- Pielke, R. A., Lyons, W. A., McNider, R. T., Moran, M. D., Moon, D. A., Stocker, R. A., Walko, R. L. and Uliasz, M. (1991) Regional and mesoscale meteorological modeling as applied to air quality studies. In *Air Pollution Modeling and Its Application VIII*, eds. H. van Dop and D. G. Steyn, pp. 259–290, Plenum Press, New York.
- Pielke, R. A. and Uliasz, M. (1993) Influence of landscape variability on atmospheric dispersion. *Journal of Air and Waste Management* **43**, 989–994.
- Portela, A. and Castro, M. (1996) Summer thermal lows in the Iberian peninsula: a three-dimensional simulation. *Quarterly Journal of the Royal Meteorological Society* **122**, 1–22.
- Ratto, C. F., Festa, R., Romeo, C., Frumento O. A. and Galluzzi, M. (1994) Mass-consistent models for wind fields over complex terrain: the state of the art. *Environment and Software* **9**, 247–268.
- Schayes, G., Thunis, P. and Bornstein, R. (1996) Topographic vorticity-model mesoscale- β (TVM) model. Part I: formulation. *Journal of Applied Meteorology*, in press.

- Schichtel, B. and Husar, R. (1994) The CAPITA Monte Carlo Model: PC implementation. In *Aerosols and Atmospheric Optics: Radiative Balance and Visual Air Quality*, Air and Waste Management Association, 26–30 September 1994, Snowbird, UT, pp. 578–600.
- Schlünzen, K. H. (1994) Mesoscale modelling in complex terrain — an overview on the German nonhydrostatic models. *Beitraege of Physics and Atmosphere* **67**, 243–253.
- Sharan, M., Gopalakrishnan, S. G., McNider, R. T. and Singh, M. P. (1996) Bhopal gas leak: a numerical investigation of the prevailing meteorological conditions. *Journal of Applied Meteorology* **35**, 1637–1657.
- Sirois, A., Olson, M. and Pabla, B. (1995) The use of spectral analysis to examine model and observed O₃ data. *Atmospheric Environment* **29**, 411–422.
- Sistla, G., Zhou, N., Hao, W., Ku, J.-Y., Rao, S. T., Bornstein, R., Freedman, F. and Thunis, P. (1996) Effects of uncertainties in meteorological inputs on Urban Airshed Model predictions and ozone control strategies. *Atmospheric Environment* **30**, 2011–2025.
- Stocker, R. A., Pielke, R. A., Verdon, A. J. and Snow, J. T. (1990) Characteristics of plume releases as depicted by balloon launchings and model simulations. *Journal of Applied Meteorology* **29**, 53–62.
- Svensson, G. (1996a) A numerical model for chemical and meteorological processes in the atmospheric boundary layer. Part I: A model description and a one-dimensional parameter study. *Journal of Applied Meteorology* **35**, 939–954.
- Svensson, G. (1996b) A numerical model for chemical and meteorological processes in the atmospheric boundary layer. Part II: A case study of the air quality situation in Athens, Greece *Journal of Applied Meteorology* **35**, 955–973.
- Tucker, G. B. (1993) New skills in predicting atmospheric pollution. *Australian Meteorological Magazine* **42**, 163–174.
- Uliasz, M. (1994) Subgrid scale parameterizations, In: *Mesoscale Modeling of the Atmosphere* (edited by Pearce, R. and Pielke, R. A.), pp. 13–19. American Meteorological Society, Boston.
- Uliasz, M., Stocker, R. A. and Pielke, R. A. (1994) Numerical modeling of air pollution transport in the southwestern United States. In *Aerosols and Atmospheric Optics: Radiative Balance and Visual Air Quality*, Air and Waste Management Association, 26–30 September 1994, Snowbird, UT, pp. 1229–1239.
- Uliasz, M., Stocker R. A. and Pielke, R. A. (1996) Regional modeling of air pollution transport in the Southwestern United States. In: *Environmental Modeling III*, Ed. P. Zannetti, Computational Mechanics Publications, Southampton, UK, pp. 145–182.
- Venkatram, A., Karamchandani, P., Pai, P. and Saxena, P. (1994) Source-receptor relationships for visibility on the Colorado Plateau. I. Modeling approach. In *Aerosols and Atmospheric Optics: Radiative Balance and Visual Air Quality*, Air and Waste Management Association, 26–30 September 1994, Snowbird, UT, pp. 693–720.
- Welander, P. (1955) studies on the general development of motion in a two-dimensional ideal fluid. *Tellus* **7**, 141–156.
- Wheeler, N. J. M. (1996) Experience based procedures for model performance evaluation. *Preprints, 9th Joint Conference on Applications of Air Pollution Meteorology with A&WMA*, 28 January–2 February 1996, Atlanta, Georgia, American Meteorological Society, Boston, pp. 512–515.
- Yamada, T. (1996) A numerical simulation of cloud distributions over coastal complex terrain. *Preprints, 9th Joint Conference on Applications of Air Pollution Meteorology with A&WMA*, 28 January–2 February 1996, Atlanta, Georgia, American Meteorological Society, Boston, pp. 498–410.
- Zannetti, P. (1990) *Air Pollution Modeling. Theories, Computational Methods and Available Software*. Computational Mechanics Publications, Southampton, UK, 444 pp.